

MEETING ABSTRACT

Open Access

Lipid metabolism does not influence the expression of proximal aortopathy in bicuspid aortic valve disease

Tatiana M Sequeira Gross*, Thomas Kuntze, Evaldas Girdauskas

From World Society of Cardiothoracic Surgeons 25th Anniversary Congress, Edinburgh
Edinburgh, UK. 19-22 September 2015

Background/Introduction

Controversy exists regarding the pathogenesis of bicuspid aortic valve (BAV) associated aortopathy. Recent data indicate the potential role of lipid metabolism in the expression of aortopathy.

Aims/Objectives

We aimed to correlate the markers of lipid metabolism with severity of proximal aortopathy in patients with BAV vs. tricuspid aortic valve (TAV) disease.

Method

A total of 458 consecutive patients (mean age 64 ± 11 years, 68% male) underwent aortic valve replacement (AVR) with/without proximal aortic surgery from January, 2008 through December, 2014. All patients undergoing combined procedures (e.g., AVR+CABG) were excluded. Only patients in whom proximal aortic dimensions were defined by preoperative CT/MRI and/or TOE were included. Correlation analysis was performed between markers of lipid metabolism (i.e., cholesterol, LDL, HDL, and triglyceride) and maximal diameter of the proximal aorta in BAV subgroup ($n = 273$) vs. TAV subgroup ($n = 185$). Moreover, we compared correlation patterns between markers of lipid metabolism and maximal aortic diameter in BAV insufficiency ($n = 46$) vs. BAV stenosis ($n = 227$) cohorts. Logistic regression was performed to identify risk factors for proximal aortic diameter >40 mm in BAV and TAV subgroups.

Results

No correlation was found between markers of lipid metabolism and proximal aortic diameter in BAV subgroup

($r = -0.1$, $p = 0.1$) and TAV subgroup ($r = 0.006$, $p = 0.9$). No significant differences in correlation patterns were found between markers of lipid metabolism and maximal aortic diameter in BAV insufficiency ($r = 0.03$, $p = 0.8$) vs. BAV stenosis ($r = -0.1$, $p = 0.1$) cohorts. Logistic regression analysis revealed triglyceride levels (HR 1.4, $p = 0.05$) and statin therapy (HR 0.4, $p = 0.03$) as predictors of proximal aortic diameter >40 mm in TAV subgroup only.

Discussion/Conclusion

Our study demonstrates no linear correlation between markers of lipid metabolism and proximal aortic diameters in a surgical cohort of BAV and TAV patients. Statin therapy and triglyceride levels influence significantly proximal aortic diameter in patients with TAV, but not with BAV disease.

Published: 16 December 2015

doi:10.1186/1749-8090-10-S1-A65

Cite this article as: Gross et al.: Lipid metabolism does not influence the expression of proximal aortopathy in bicuspid aortic valve disease. *Journal of Cardiothoracic Surgery* 2015 **10**(Suppl 1):A65.

Department of Cardiac Surgery, Central Clinic Bad Berka, Germany