

ORAL PRESENTATION

Open Access

Mid-term impact of patient-prosthesis mismatch on young and middle-aged patients undergoing surgery due to severe aortic stenosis

D Hernandez-Vaguero^{1*}, R Diaz¹, JM Garcia², D Calvo², JC Llosa¹

From 23rd World Congress of the World Society of Cardio-Thoracic Surgeons Split, Croatia. 12-15 September 2013

Background

The clinical impact of patient-prosthesis mismatch on outcomes in young and middle-aged patients undergoing surgery for aortic valve replacement remains unknown. Our objective was to examine the mid-term impact of patient-prosthesis mismatch on overall mortality, quality of life and cardiac events.

Methods

All patients younger than 70 years of age undergoing isolated aortic valve replacement from October 2005 to October 2011 were analyzed. Patient-prosthesis mismatch was defined as the indexed effective orifice area ≤ 0,85cm2/m2. We studied the impact of patient-prosthesis mismatch on mid-term overall mortality, cardiac events and New York Heart Association functional class using an analysis stratified for propensity score. Cardiac events were defined as cardiac death, sudden death, hospital readmission due to angina, syncope or heart failure or reoperation on aortic prosthesis.

Results

293 patients were included in the study, of whom 81 (27,61%) had some degree of patient-prosthesis mismatch. Patient-prosthesis mismatch had no impact on mid-term overall mortality (HR=1,45 CI 95%= 0,65-3,22; p=0,36), although it had a negative impact on cardiac events (HR=11,52 CI 95%=5,25-25,24; p<0,001) and functional class (RR=7,55 CI 95%=2,59-22,03; p<0,001).

Full list of author information is available at the end of the article

Conclusions

Patient-prosthesis mismatch appears to be a strong and independent predictor of cardiac events and functional class in young and middle aged patients undergoing aortic valve replacement for severe stenosis. However, it is possible that it has no impact on overall mortality.

Authors' details

¹Cardiac Surgery Department, Central University Hospital of Asturias, Oviedo, Spain. ²Department of Cardiology, Central University Hospital of Asturias, Oviedo, Spain.

Published: 11 September 2013

doi:10.1186/1749-8090-8-S1-O101

Cite this article as: Hernandez-Vaquero *et al.*: Mid-term impact of patient-prosthesis mismatch on young and middle-aged patients undergoing surgery due to severe aortic stenosis. *Journal of Cardiothoracic Surgery* 2013 8(Suppl 1):O101.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit





^{*} Correspondence: dhvaquero@gmail.com

¹Cardiac Surgery Department, Central University Hospital of Asturias, Oviedo, Spain