

POSTER PRESENTATION

Open Access

Left ventricular pseudoaneurysm after ascending, arch and descending aortic replacement with Evita Open Plus[®] prosthesis

JR Echevarría*, M Fernández, M Blanco, H Valenzuela, G Laguna, P Pelaez

From 23rd World Congress of the World Society of Cardio-Thoracic Surgeons
Split, Croatia. 12-15 September 2013

Background

We report a rare complication after successful Evita Open Plus[®] implantation with cardiopulmonary bypass and circulatory arrest complete for total aortic aneurysm treatment.

Methods

A 71 year old male with aortic aneurysm was referred for consideration of surgery. The patient comes to the emergency department to present an acute left chest pain. He has a history of duodenal ulcer and arterial hypertension. Rx thorax demonstrated mediastinal widening compatible with giant aortic aneurysm. The echocardiogram (TTE) confirmed normal sized cardiac cavities and ascending aortic aneurysm. Aortography presented severe dilatation of ascending thoracic aorta with moderate aortic regurgitation. CT Angiography established supra-aortic ascending aneurysm with maximum dimension of 53 mm, 65 mm arch aneurysm and descending 106 mm in the transverse plane.

Upper medial re sternotomy, cardiopulmonary bypass and circulatory arrest complete with moderate hypothermia were performed as well as ascending aorta and arch replacement with reimplantation of the supra-aortic trunks by hybrid prosthetic Evita Open Plus[®] 33.

Results

The patient evolution was favourable. In the control TTE was observed on the left ventricular apex two interconnected cavities support with left ventricular pseudoaneurysm, confirmed by CT angiogram. The patient required a new intervention, performing

exclusion of the pseudoaneurysm in the left ventricular (LV) apex with a Teflon patch. After surgery remains stable and was discharged without complications.

Conclusion

The total aortic replacement (ascending aorta, arch and descending aorta) using hybrid routines are reserved for patients with complex pathology. Iatrogenic LV pseudoaneurysms postoperative may uncommon. In some cases is related to the increase in consistency (due to hypothermia during circulatory arrest complete) of the cannula vent of the LV during extracorporeal circulation.

Published: 11 September 2013

doi:10.1186/1749-8090-8-S1-P2

Cite this article as: Echevarría et al.: Left ventricular pseudoaneurysm after ascending, arch and descending aortic replacement with Evita Open Plus[®] prosthesis. *Journal of Cardiothoracic Surgery* 2013 **8**(Suppl 1):P2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: joraech@hotmail.com
Hospital Clínico, Valladolid, Spain