

RESEARCH ARTICLE

Open Access

Risk factors for postoperative hypoxemia in patients undergoing Stanford A aortic dissection surgery

Yinghua Wang*, Song Xue and Hongsheng Zhu

Abstract

Background: The purpose of this study is to identify the risk factors for postoperative hypoxemia in patients with Stanford A aortic dissection surgery and their relation to clinical outcomes.

Methods: Clinical records of 186 patients with postoperative hypoxemia in Stanford A aortic dissection were analyzed retrospectively. The patients were divided into two groups by postoperative oxygen fraction (PaO₂/FiO₂): hypoxemia group (N=92) and non-hypoxemia group (N=94).

Results: We found that the incidence of postoperative hypoxemia was 49.5%. Statistical analysis by t-test and χ^2 indicated that acute onset of the aortic dissection ($p=0.000$), preoperative oxygen fraction (PaO₂/FiO₂) ≤ 200 mmHg ($p=0.000$), body mass index ($p=0.008$), circulatory arrest (CA) time ($p=0.000$) and transfusion more than 3000 ml ($p=0.000$) were significantly associated with postoperative hypoxemia. Multiple logistic regression analysis showed that preoperative hypoxemia, CA time and transfusion more than 3000 ml were independently associated with postoperative hypoxemia in Stanford A aortic dissection.

Conclusion: Our results suggest that postoperative hypoxemia is a common complication in patients treated by Stanford A aortic dissection surgery. Preoperative oxygen fraction lower than 200 mmHg, longer CA time and transfusion more than 3000 ml are predictors of postoperative hypoxemia in Stanford A aortic dissection.

Background

Aortic dissection can be divided into type A and type B by Stanford classification and is one of the most urgent medical cases for its high mortality and morbidity. Although open heart surgery for treating Stanford A aortic dissection has been a universal option, the patients are still threatened by the lethal perioperative complications such as hemorrhage, hypoxemia, renal dysfunction, cerebral dysfunction and etc. Postoperative hypoxemia is defined as an oxygen fraction (PaO₂/FiO₂) ≤ 200 mmHg under mechanical ventilation in the first 24 hour after surgery [1]. As a matter of fact, postoperative hypoxemia resembles ARDS (acute respiratory distress syndrome) when oxygen fraction is below 200 mmHg.

Patients who undergo cardiac surgery generally have a more or less increased risk of postoperative hypoxemia. Postoperative hypoxemia has been reported to occur in

12.2-27.1% of patients after cardiopulmonary bypass, and in as high as 51% after open heart aortic dissection surgery [1-3]. Hypoxemia is a dangerous complication and usually accompanied by a number of untoward consequences such as prolonged ventilator support, longer stay in ICU and hospital. Prolonged mechanical ventilation would lead to more and serious complications. Continuous postoperative hypoxemia may cause dysfunctions of other organs besides lungs. The aim of our study is to evaluate the risk factors of postoperative hypoxemia in Stanford A aortic dissection and its impact on the clinical outcomes.

Methods

From Dec 2004 to July 2012, 186 consecutive patients with acute Stanford A aortic dissection underwent open heart surgery in RenJi Hospital. This study was approved by RenJi Hospital Clinical Research Ethics Committee (2012027). Acute Stanford A aortic dissection was diagnosed by enhanced computed tomography scan and aortic valve regurgitation was confirmed by echocardiography.

* Correspondence: pen1019@hotmail.com
Cardiovascular Surgery Department, RenJi Hospital, School of medicine,
Shanghai JiaoTong University, 1630 Dongfang Rd., Shanghai 200127, China

About 70% of the patients underwent surgical procedure within 48 hours, others within 21 days. 4 patients who died during the first 48 hours after surgery were the result of multiorgan dysfunction syndrome and refractory bleeding. They have been excluded from the 186 patients above. In the present study, namely postoperative hypoxemia was defined as a PaO₂/FiO₂ ≤200 mmHg in the first 24 hour after surgery, according to the diagnostic criteria for ARDS established by American-European consensus conference. The 186 patients were divided into two groups by postoperative oxygen fraction (PaO₂/FiO₂): hypoxemia group (N=92) and non-hypoxemia group (N=94). We compared 17 perioperative factors between the two groups. Mortality, morbidity and long-term survival were compared between the two groups. In-hospital outcome variables were obtained from patients' medical records. Follow-up information including survival, cardiovascular-related events and causes of death were obtained through out-patient clinic visit or from telephone correspondence with the patients or their relatives.

Surgical procedure

All operations were performed through a median sternotomy. After heparinization, cardiopulmonary bypass was established by cannulation of right atrium or separately the superior and inferior vena cava. The right or left femoral artery or the right subclavian artery was optionally the site of cannulation. A left ventricular vent line was inserted through the right superior pulmonary vein. Once the cardiopulmonary bypass started, systemic cooling was initiated. After clamping the ascending aorta, the root of aorta was opened and cardioplegic solution was infused through direct selective cannulation of the coronary ostia and the intimal tear and entry of the dissection was detected. When the proximal aortic root procedure was finished, the rectal temperature was around 18°C. The circulation was arrested, crossclamping removed and the intimal tear or entry of dissection of aortic arch and proximal descending aorta was explored and identified. If the entry was close to the orifice of the arch vessels, aortic arch was totally replaced, and if the entry was localized in the lesser curvature of the transverse aortic arch proximal to the left subclavian artery, hemiarch replacement or intraoperative descending aorta endograft deployment was performed. When the intimal tear involved the ascending aorta and the entry site could not be identified, the entire ascending aorta was replaced. Aortic root replacement, Bentall procedure would be performed for patients with dilation of aortic root, especially in Marfan's syndrome patients. Antegrade selective cerebral perfusion was carried out when we expected that the circulatory arrest time would be longer than 30 min or determined by the surgeon's personal preference and experience. With the help of antegrade selective cerebral

perfusion, the rectal temperature may be maintained at about 25°C during circulatory arrest. The operative variables and concomitant procedures are shown in Table 1. Arch-involved procedure was performed in 84 patients,

Table 1 Operative variables and concomitant procedures

Operative data	Total (n=186)
Arterial cannulation	
Femoral artery	178
Right subclavical artery	8
Venous canulation	
Bicaval	6
Right atrium	180
Perfusion technique	
ECC+DHCA	178
ECC+ASCP	8
Operative procedures	
Arch-involved procedure	84
Total arch replacement (TAR)	48
Hemiarch replacement (HAR)	14
Intraoperative aortic arch endograft deployment (IAAED)	22
Intraoperative descending aorta endograft deployment (IDAED)	66
Wheat procedure	6
Wheat+ TAR+IDAED	2
Bentall's procedure	34
Bentall's +HAR	4
Bentall's +IAAED	4
Bentall's +TAR+IDAED	8
Bentall's +IDAED	2
Ascending aorta replacement (AAR)	40
AAR+IAAED	18
AAR+TAR	4
AAR+HAR	10
AAR+IDAED	20
AAR+TAR+IDAED	34
Concomitant procedures	
Cornary artery bypass grafting	4
Mitral valve replacement	4
Atrial septal defect repair	2
CPB time	184.85±36.83
AO time	94.10±34.58
ASCP time	50.63±2.39
CA time	26.19±9.91

ECC:extracorporeal circulation; ASCP:antegrade selective cerebral perfusion; IAAED:Intraoperative aortic arch endograft deployment; IDAED:Intraoperative descending aorta endograft deployment; CPB:cardiopulmonary bypass; AO: aortic clamping; CA:circulatory arrest.

including 48 total arch replacement, 14 hemiarch replacement and 22 intraoperative aortic arch endograft deployment. Concomitant procedures included coronary artery bypass grafting in 4 patients, mitral valve replacement in 4 patients and atrial septal defect repair in another 2 patients.

Statistical analysis

All statistical analysis were performed with SPSS16.0 for windows. Quantitative variables were described as mean and standard deviation. The test of homogeneity variance was used for determination of quantitative data distribution. When the distribution of variables was normal, independent samples t-test was used for comparison of quantitative sizes of two independent samples. The dependence of qualitative variable was evaluated by Chi-squared (χ^2) criterion. Multivariate stepwise forward logistic regression analysis was adopted to identify independent risk factors for postoperative hypoxemia. Survival was analyzed by the Kaplan-Meier method and log rank test. A p value of less than 0.05 was considered statistically significant. Kaplan-Meier curves were shown in Figure 1.

Results

From Dec 2004 to July 2012, 186 patients with acute Stanford A aortic dissection underwent open heart surgery in RenJi Hospital. The age of the patients was 51.42 ± 8.93 years, ranging from 28 to 76 years. 70.97% of them were male and 78.5% were diagnosed as acute aortic dissection. The overall in-hospital mortality was 1.1%. The perioperative variables of patients were shown in Table 2.

As shown in Table 2, the incidence of postoperative hypoxemia in our patient population was 49.5%. In order

Table 2 Preoperative patient characteristics

Characteristics	Total (n=186)
Age (years), M \pm SD	51.42 \pm 8.93
Gender (male%)	70.97
Height (cm), M \pm SD	169.19 \pm 8.83
Weight (kg), M \pm SD	70.94 \pm 15.23
BMI (kg/m ²), M \pm SD	24.68 \pm 4.48
History of smoking (%)	32.8
Acute onset (%)	78.49
Preoperative hypoxemia (%)	49.46
History of previous cardiac operation (%)	4.30
Preoperative LVEF, M \pm SD	55.54 \pm 7.50
Hypertension (%)	57.5
Diabetes mellitus (%)	5.38

BMI-body mass index, LVEF-left ventricular ejection fraction.
M \pm SD-mean \pm standard deviation.

to determine the factors associated with postoperative hypoxemia, 186 patients were divided into two groups by their postoperative oxygen fraction (PaO₂/FiO₂). Patients with postoperative PaO₂/FiO₂ \leq 200 mmHg were classified into the hypoxemia group, while others into the non-hypoxemia group. Univariate analyses of perioperative characteristics in these two groups were shown in Table 3. Significant differences between the two groups were found in following clinical variables: acute onset of the aortic dissection (p=0.000), preoperative oxygen fraction (PaO₂/FiO₂) \leq 200 mmHg (p=0.000), body mass index (p=0.008), circulatory arrest (CA) time (p=0.000) and transfusion more than 3000 ml (p=0.000).

As shown in Table 4, multiple logistic regression analysis confirmed that preoperative hypoxemia, CA time




Table 3 Univariate analysis in postoperative hypoxemia

Characteristics	Hypoxemia group	Non-hypoxemia group	P value
Age (years), mean (sd)	51.70 (9.16)	51.15 (8.74)	NS
Gender (male) (%)	50	50	NS
Height (cm), mean (sd)	168.26 (8.48)	170.11 (9.10)	NS
Weight (cm), mean (sd)	72.48 (15.16)	69.44 (15.22)	NS
BMI (kg/m ²), mean (sd)	25.55 (4.69)	23.83 (4.12)	P=0.008
History of smoking (%)	57.4	42.6	NS
Acute onset (%)	56.2	43.8	P=0.000
Preoperative hypoxemia (%)	76.1	23.9	P=0.000
History of previous cardiac operation (%)	25	75	NS
Preoperative LVEF (%), mean (sd)	56.50 (5.55)	54.60 (8.94)	NS
Hypertension (%)	53.3	46.7	NS
Diabetes mellitus (%)	80	20	NS
CPB time (min), mean (sd)	185.50 (37.61)	184.21 (36.24)	NS
AO time (min), mean (sd)	94.86 (33.97)	93.35 (35.33)	NS
CA time (min), mean (sd)	29.29 (10.29)	23.17 (8.54)	P=0.000
Postoperative LVEF<40% (%)	66.7	33.3	NS
mechanical ventilation duration (hour), mean (sd)	79.22 (62.31)	41.45 (5.54)	NS
Postoperative transfusion ≥3000 ml (%)	76.1	23.9	P=0.000

BMI-body mass index, LVEF-left ventricular ejection fraction.
 CPB-cardiopulmonary bypass AO-aortic clamping.
 CA- circulatory arrest.
 m(sd)—mean(standard deviation).

and transfusion more than 3000 ml were independently associated with postoperative hypoxemia in Stanford A aortic dissection.

All patients have been followed up through out-patient clinic visit or by telephone contacts. The follow-up period was from one month to ninety months (43.65±23.70 month), with a period of 47.96±24.10 month for hypoxemia group and 39.44±22.64 month for non-hypoxemia group. Four patients died, two died of respiratory failure during the first three month after operation in the hospital, others died of stroke or sudden death during follow-up period. Kaplan-Meier survival curve was shown in Figure 1 which

revealed a significant difference in the 90-month survival between the hypoxemia group and non-hypoxemia group (p=0.014).

Discussion

Stanford A aortic dissection has been known for its high mortality and morbidity. Since deep hypothermia, ultrafiltration and selective cerebral perfusion technique were applied in surgery for Stanford A aortic dissection, the incidence of postoperative cerebral complications has decreased. Hypoxemia, as another life-threatening postoperative complication for Stanford A aortic dissection, may

Table 4 Multivariate logistic regression analysis model for postoperative hypoxemia

		Variables in the equation								
		B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)		
									Lower	Upper
Step 1 ^a	BMI	.020	.050	.151	1	.697	1.020	.924	1.125	
	Acute onset	-.859	.626	1.886	1	.170	.424	.124	1.444	
	Preoperative hypoxemia	2.682	.528	25.758	1	.000	14.611	5.187	41.158	
	CA time	.078	.023	11.286	1	.001	1.081	1.033	1.132	
	transfusion≥3000 ml	2.256	.425	28.211	1	.000	9.541	4.151	21.932	
	Constant	-4.368	1.386	9.924	1	.002	.013			

a. Variable(s) entered on step 1: BMI, acute onset of aortic dissection, preoperative hypoxemia, CA time, transfusion ≥3000 ml.
 BMI-body mass index.
 CA- circulatory arrest.

be attributed to prolonged mechanical ventilation and ICU-stay. Acute aortic dissection is frequently accompanied by hypoxemia. Yet currently, few studies revealed the clinical entity in such patients and the management after surgery for acute Stanford A aortic dissection surgery. The aim of our study is to reveal the incidence, risk factors and management of hypoxemia after the surgery for acute Stanford A aortic dissection.

Previous studies disclosed that persistent hypoxemia was the most common cause of PMV (prolonged mechanical ventilation) and ARDS was the most common cause of hypoxemia [4-6]. The etiology of acute respiratory distress syndrome and its less severe form acute lung injury (ALI) may be systemic, rather than pulmonary confirmed to. One of the diagnostic criteria of ARDS is oxygenation impairment ($\text{PaO}_2/\text{FiO}_2 \leq 200$ mmHg). Nevertheless, the respiratory failure in patients with hypoxemia in this study could not be diagnosed technically as ARDS or ALI, because it was not usually accompanied by typical radiological pulmonary infiltrates. Therefore, in the present study, postoperative hypoxemia was defined as $\text{PaO}_2/\text{FiO}_2 \leq 200$ mmHg in the first 24 hour after surgery, according to the diagnostic criteria for ARDS established by American-European consensus conference [4-8].

Patients who undergo cardiac surgery have an increased risk of postoperative hypoxemia [5]. Postoperative hypoxemia has been reported to occur in 12.2%-27.1% of patients after cardiopulmonary bypass and usually is 51% in open heart aortic dissection surgery [1-3]. In this study, the incidence of postoperative hypoxemia is 49.5% in open heart Stanford A dissection surgery. Previously reported risk factors for hypoxemia after cardiac surgery include advanced age, obesity, smoking history, previous heart surgery, emergency surgery, low LVEF, preoperative chronic pulmonary disease, preoperative myocardial infarction, preoperative diabetes, cardiogenic and noncardiogenic pulmonary edema, pneumonia, excessive blood transfusion and prolonged CPB time [9-11]. In the present study, we identified BMI, preoperative hypoxemia ($\text{PaO}_2/\text{FiO}_2 \leq 200$ mmHg), acute onset of Stanford A aortic dissection, CA time, postoperative transfusion ≥ 3000 ml as the risk factors for postoperative hypoxemia after Stanford A aortic dissection surgery.

Postoperative hypoxemia is a common complication in cardiac surgery. CPB (cardiopulmonary bypass) and CA (circulatory arrest) have been reported to play important roles in the postoperative hypoxemia [12-16]. Stanford A aortic dissection surgery is associated with a high incidence of postoperative hypoxemia compared with other elective cardiac surgical procedures [5,14-16]. During cardiac surgery, both experimental and clinical studies have well documented that open heart surgery with CPB could result in systemic inflammatory response, activation

of complements, thrombin, cytokines, endothelin, endotoxins, neutrophils, adhesion molecule, macrophages, multiple inflammatory mediators and impaired immune reaction and organ dysfunction [17-23]. As CPB represents a kind of non-physiological circulation affecting the peripheral tissue perfusion, particularly in the case of prolonged pump time, which might lead to impaired capillary membrane permeability, malperfusion, tissue anoxia and pulmonary complications [24-27]. Although deep hypothermia and antegrade selective cerebral perfusion may preserve better cerebral function during circulatory arrest, the malperfusion of other organs may still lead to generally ischemic reperfusion injury. A prospective study by De Santo et al. analysed the effect of pulmonary perfusion during hypothermic circulatory arrest which may preserve better lung function but that was not used in our study [28].

Deep hypothermia is an important technique commonly used in acute Stanford A aortic dissection surgery. However, deep hypothermia participates in both platelet activation pathways and the enzymatic activity of clotting factors which may lead to bleeding and subsequent transfusion. Excessive transfusion is attributed to active bleeding or coagulation abnormalities. Loss of coagulation factors, destruction of platelets and microthrombosis may all lead to lung injury [29-31]. In our study, prolonged CA time and transfusion more than 3000 ml are associated with the hypoxemia.

According to our experience, large dose of methylprednisolone in the operation and a rather small dose of methylprednisolone during the first three day after the operation are regularly used in order to decrease the inflammatory reaction. We also run the operation and control bleeding as fast as possible to reduce the CPB or DHCA (deep hypothermic circulatory arrest) time and excessive transfusion.

In the present study, 78.49% of patients who were sent to emergency department for acute onset of Stanford A aortic dissection were found to be hypoxemia ($\text{PaO}_2/\text{FiO}_2 \leq 200$ mmHg) before the operation. In our study, all patients breathed spontaneously with nasal prongs or face mask with inhaled oxygen 5-8 L/min before operation. The arterial blood gas and preoperative $\text{PaO}_2/\text{FiO}_2$ were calculated before mechanical ventilation. Both acute onset of dissection and preoperative hypoxemia are associated with postoperative hypoxemia. Acute onset of aortic dissection indicates sudden rupture of intima, propagation of the dissection into the medial layer, acute bleeding and activation of cellular and humoral inflammatory systems. In the process of acute bleeding, ventilation and perfusion are out of balance, which leads to hypoxemia. Quite a number of investigators reported that oxygenation was impaired from the onset of dissection due to the inflammatory cascade reaction, increased

endothelial and epithelial permeability, increased pulmonary vascular pressure and impaired alveolar surfactant function, which can lead to decrease in the preoperative PaO₂/FiO₂ ratio. The preoperative inflammatory status of the patient significantly affects the degree of systemic inflammatory response during the Stanford A aortic dissection surgery in succession [1,9]. Therefore, postoperative hypoxemia may be related to preoperative disturbance in oxygenation. In our opinion, we try all the best to maintain the perioperative fluid balance and meticulous ventilation manipulation, especially in the very first day after the operation.

Obesity have been reported as predictors for hypoxemia [4,32]. Consistently, we found that obese patients, with BMI > 25 kg/m² are more likely to develop postoperative hypoxemia. It is well-known that obesity is frequently accompanied by some co-morbidity, including obstructive sleep apnea and hypoventilation syndrome which all related to hypoxia [32,33]. In the present study, the body mass index showed a significant influence in univariate analysis, but not in the multivariate regression model.

In our study, the long-term survival in non-hypoxemia group was better than that in hypoxemia group, which was in accordance with previous studies [13]. Survivors following cardiac surgery are found to enjoy a good quality of life after discharge.

Conclusions

In conclusion, our study demonstrated that BMI (Body mass index), preoperative hypoxemia (PaO₂/FiO₂ ≤ 200 mmHg), acute onset of Stanford A aortic dissection, CA time, postoperative transfusion ≥ 3000 ml were the key risk factors for Stanford A aortic dissection surgery. And preoperative hypoxemia, CA time and transfusion ≥ 3000 ml were independent predictors of postoperative hypoxemia after Stanford A aortic dissection surgery. Optimized treatments including preoperative respiratory tract management to control those predictors should be helpful to improve the clinical outcomes.

Limitation

Like all retrospective studies, this investigation also has several limitations. First, the patient population collected for investigation was relatively small. Second, owing to the emergency nature of the Stanford A aortic dissection surgery, detailed information about preoperative pulmonary disease was not completely available. The presence of COPD (chronic obstructive pulmonary disease) was judged on the basis of the patient's medical history which might not be well defined. Finally, as this is a retrospective study, the potential misclassification bias could not be completely excluded.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SX performed all the surgeries in our study. HZ helped to design the study and revise the grammar and idiomatic expression. YW participated in design of the study, statistically analysis and draft of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We acknowledge the Dr. Weijun Wang from RenJi Hospital for his help in acquisition of data. Tao Zhang from Ruijin Hospital for her assistance with statistical analysis.

Received: 4 November 2012 Accepted: 29 April 2013

Published: 30 April 2013

References

1. Takayuki N, Kohei K, Hiroshi I, Tsuyoshi K, Hiroyuki N, Nobuo S: **Risk factors for hypoxemia after surgery for acute type A aortic dissection.** *Surg Today* 2006, **36**:680–685.
2. Filsoufi F, Rahmanian PB, Castillo JG, Chikwe J, Adams DH: **Logistic risk model predicting postoperative respiratory failure in patients undergoing valve surgery.** *Eur J Cardiothorac Surg* 2008, **34**(5):953–959.
3. Ji Q, Mei Y, Wang X, Feng J, Cai J, Sun Y, Xie S, Li D, Dayi H: **Study on the risk factors of postoperative hypoxemia in patients undergoing coronary artery bypass grafting.** *Circ J* 2008, **72**:1975–1980.
4. Yende S, Wunderink R: **Causes of prolonged mechanical ventilation after coronary artery bypass surgery.** *Chest* 2002, **122**:245–252.
5. Branca P, McGaw P, Light RW: **Factors associated with prolonged mechanical ventilation following coronary artery bypass surgery.** *Chest* 2001, **119**:537–546.
6. Filsoufi F, Rahmanian PB, Castillo JG, Chikwe J, Adams DH: **Predictors and early and late outcomes of respiratory failure in contemporary cardiac surgery.** *Chest* 2008, **133**(3):713–721.
7. Girdeauskas E, Kuntze T, Borger MA, et al: **Acute respiratory dysfunction after surgery for acute type A aortic dissection.** *Eur J Cardiothorac Surg* 2010, **37**:691–696.
8. Kurabayashi M, Okishige K, Azegami K, et al: **Reduction of the PaO₂/FiO₂ ratio in acute aortic dissection.** *Circ J* 2010, **74**:2066–2073.
9. Weiss YG, Merlin G, Koganov E, Ribo A, Oppenheim-Eden A, Medalion B, et al: **Postcardiopulmonary bypass hypoxemia: a prospective study on incidence, risk factors and clinical significance.** *J Cardiothorac Vasc Anesth* 2001, **14**:506–513.
10. Milot J, Perron J, Lacasse Y, Letourneau L, Cartier PC, Maltais F: **Incidence and predictors of ARDS after cardiac surgery.** *Chest* 2001, **119**:884–888.
11. Szeles TF, Yoshinaga EM, Alenca W, Brudniewski M, Ferreira FS, Auler JO, Carmona MJ, Malbouisson LM: **Hypoxemia after myocardial revascularization: analysis of risk factors.** *Rev Bras Anestesiol* 2008, **58**(2):124–136.
12. Salis S, Mazzanti VV, Merli G, Salvi L, Tedesco CC, Veglia F, Sisillo E: **Cardiopulmonary bypass duration is an independent predictor of morbidity and mortality after cardiac surgery.** *J Cardiothorac Vasc Anesth* 2008, **22**(6):814–822.
13. Kimura N, Tanaka M, Kawahito K, Sanui M, Yamaguchi A, Ino T, Adachi H: **Risk factors for prolonged mechanical ventilation following surgery for acute type A aortic dissection.** *Circ J* 2008, **72**:1751–1757.
14. Ip P, Chiu CS, Cheung Y: **Risk factors prolonging ventilation in young children after cardiac surgery: impact of noninfectious pulmonary complication.** *Pediatr Crit Care Med* 2002, **3**:269–274.
15. Garcia Montes JA, Calderon Colmenero J, Casanova M, et al: **Risk factors for prolonged mechanical ventilation after surgical repair of congenital heart disease.** *Arch Cardiol Mex* 2005, **75**:402–407.
16. Szekely A, Sapi E, Kiraly L, et al: **Intraoperative and postoperative risk factors for prolonged mechanical ventilation after pediatric cardiac surgery.** *Paediatr Anaesth* 2006, **16**:1166–1175.
17. Chiappini B, Schepens M, Tan E, Dell Amore A, Morshuis W, Dossche K, et al: **Early outcomes of acute type A aortic dissection: analysis of risk factors in 487 consecutive patients.** *Eur Heart J* 2004, **26**:180–186.
18. Sugano Y, Anzai T, Yoshikawa T, Satoh T, Iwanaga S, Hayashi T, et al: **Serum C-reactive protein elevation predicts poor clinical outcome in patients**

- with distal type acute aortic dissection: association with the occurrence of oxygenation impairment. *Int J Cardiol* 2005, **102**:39–45.
19. Kallenbach K, Oelze T, Salcher R, Hagl C, Karck M, Leyh RG, *et al*: Evolving strategies for treatment of acute aortic dissection type A. *Circulation* 2004, **110**(Suppl 1):II243–II249.
 20. Shi S, Zao Z, Liu X, Shu Q, Linhua Tan R, Lin ZS, Fang X: Perioperative risk factors for prolonged mechanical ventilation following cardiac surgery in neonates and young infants. *Chest* 2008, **134**:768–774.
 21. Cislaghi F, Condemi AM, Corona A: Predictors of prolonged mechanical ventilation in a cohort of 3269 CABG patients. *Minerva Anestesiol* 2007, **73**:615–621.
 22. Kin H, Kawazoe K, Nakajima K, Niinuma T, Kataoka T, Endo S, *et al*: Perioperative serum procalcitonin concentrations in patients with acute aortic dissection. *Eur Surg Res* 2003, **35**:451–454.
 23. Hasegawa Y, Ishikawa S, Ohtaki A, Otani Y, Takahashi T, Sato Y, *et al*: Impaired lung oxygenation in acute aortic dissection. *J Cardiovasc Surg (Torino)* 1999, **40**:191–195.
 24. Rothenburger M, Soeparwata R, Deng MC, Schmid C, Berendes E, Tjan TD, *et al*: Prediction of clinical outcome after cardiac surgery: the role of cytokines, endotoxin and anti-endotoxin core antibodies. *Shock* 2001, **16**(Suppl 1):44–50.
 25. Raja SG, Dreyfus GD: Modulation of systemic inflammatory response after cardiac surgery. *Asian Cardiovasc Thorac Ann* 2005, **13**:382–395.
 26. Baufreton C, Corbeau JJ, Pinaud F: Inflammatory response and haematological disorders in cardiac surgery: toward a more physiological cardiopulmonary bypass. *Ann Fr Anesth Reanim* 2006, **25**:510–520.
 27. Asimakopoulos G: Systemic inflammation and cardiac surgery an update. *Perfusion* 2001, **16**:353–360.
 28. De Santo LS, Romano G, Amarelli C, *et al*: Surgical repair of acute type A aortic dissection: continuous pulmonary perfusion during retrograde cerebral perfusion prevents lung injury in a pilot study. *J Thorac Cardiovasc Surg* 2003, **126**:826–831.
 29. Khan H, Belsher J, Yilmaz M, Afessa B, Winters JL, Moore SB, Hubmayr RD, Gajic O: Fresh-frozen plasma and platelet transfusions are associated with development of acute lung injury in critically ill medical patients. *Chest* 2007, **131**(5):1308–1314.
 30. Zilberberg MD, Carter C, Lefebvre P, Raut M, Vekeman F, Duh MS, Shorr AF: Red blood cell transfusions and the risk of acute respiratory distress syndrome among the critically ill: a cohort study. *Crit Care* 2007, **11**(3):R63.
 31. Székely A, Cserép Z, Sági E, Breuer T, Nagy CA, Vargha P, Hartyánszky I, Szatmári A, Treszl A: Risks and predictors of blood transfusion in pediatric patients undergoing open heart operations. *Ann Thorac Surg* 2009, **87**(1):187–197.
 32. Lumachi F, Marzano B, Fanti G, Basso SM, Mazza F, Chiara GB: Relationship between body mass index, age and hypoxemia in patients with extremely severe obesity undergoing bariatric surgery. *In Vivo* 2010, **24**(5):775–777.
 33. Gong MN, Thompson BT, Williams P, Pothier L, Boyce PD, Christiani DC: Clinical predictors of and mortality in acute respiratory distress syndrome: potential role of red cell transfusion. *Crit Care Med* 2005, **33**(6):1191–1198.

doi:10.1186/1749-8090-8-118

Cite this article as: Wang *et al*: Risk factors for postoperative hypoxemia in patients undergoing Stanford A aortic dissection surgery. *Journal of Cardiothoracic Surgery* 2013 **8**:118.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

